

Exactly one head: $E = \{HT, TH\}$ $P(E) = \frac{2}{4}$

At least one head: $E = \{HT, TH, HH\}$ $P(E) = \frac{3}{4}$

Exactly two heads: $E = \{HH\}$ $P(E) = \frac{1}{4}$

Exactly three heads: $E = \emptyset$ $P(E) = \frac{0}{4}$

Our main tools for these problems are the counting techniques from the last chapter

Ex 2: What is the probability that when two dice are rolled that their sum is exactly 7?

Sample space:

$$S = \{\text{all pairs of rolls}\}$$

$$= \{(1,1), (1,2), \dots, (2,1), (2,2), \dots, (6,6)\}$$

6 possibilities for each roll

So by the product rule,

$$|S| = 6 \cdot 6 = 36$$

Event:

$$E = \{\text{rolls that add to 7}\}$$

$$= \{(1,6), (2,5), (3,4), (4,3), (5,2), (6,1)\}$$

$$|E| = 6$$

$$\text{So } P(E) = \frac{|E|}{|S|} = \frac{6}{36}$$

Ex 6: What is the probability that a poker hand contains 3 of one kind, 2 of another kind (full house)?

Sol'n: Sample space: $S = \{5\text{-card hands}\}$

$$|S| = \binom{52}{5} = \frac{52!}{5!47!}$$

Event: $E = \{\text{Hands w/ full house}\}$

To choose an element of E :

- Choose the kind for the 3-of-a-kind: $\binom{13}{1}$ ways
- Choose 3 cards of this kind: $\binom{4}{3}$ ways
- Choose the kind for the pair: $\binom{12}{1}$ ways
- Choose 2 cards of this kind: $\binom{4}{2}$ ways

By prod. rule,

$$|E| = \binom{13}{1} \binom{4}{3} \binom{12}{1} \binom{4}{2} = 13 \cdot 4 \cdot 12 \cdot 6 = 3744$$

$$P(E) = \frac{|E|}{|S|} = \frac{3744}{2598960} = 0.0014$$

§ 7.2: Probability Theory

Now we make one small change. Instead of every outcome s being equally likely, they now have an individual prob. $p(s)$

$$0 \leq p(s) \leq 1 \quad \text{for all } s \in S$$

$$\sum_{s \in S} p(s) = 1$$

Note: $p: S \rightarrow [0,1]$ is a function, called a probability distribution.

Everything we do in this section also holds for the equally-likely situation, which is just $p(s) = \frac{1}{|S|}$ for all $s \in S$
"uniform distribution"

Ex 2: Roll a die where 3 is twice as likely to be rolled as any other number (which are equally likely)

Class activity: Find the prob. of rolling

- a) An odd number
- b) An even number
- c) An even number or a 3
- d) A number 3 or less
- e) An odd number or a number 3 or less

Recall the complement $\bar{E} = S \setminus E$ of E

- 1) Complement rule: $p(\bar{E}) = 1 - p(E)$
- 2) Subtraction rule: $p(E_1 \cup E_2) = p(E_1) + p(E_2) - p(E_1 \cap E_2)$
- 3) Sum rule: if E_1 & E_2 disjoint, $p(E_1 \cup E_2) = p(E_1) + p(E_2)$