

Announcements

Reminder: HW8 due Wed. 4/8

Class next Friday (4/10) cancelled

Recall: A relation from A to B is a subset of $A \times B$.
(a relation where each $a \in A$ appears exactly once is a function)

Properties:

- R is reflexive if $a R a$ for all $a \in A$
- R is symmetric if whenever $a R b$, then $b R a$
- R is antisymmetric if whenever $a R b$ and $a \neq b$, then $b \not R a$
- R is transitive if whenever $a R b$ and $b R c$, then $a R c$

Def: An equivalence relation on A is a rel'n on A which is reflexive, symmetric, and transitive

" a is equiv. to itself"

"if a is equiv. to b , then b is equiv. to a "

"if a and b are equiv., and b and c are equiv., then a and c are equiv."

Often write $a \sim b$ for "a is equiv. to b"

Def: The (maximal) subsets of A whose elts. are all equiv. are called the equivalence classes of A .

If $a \in A$, $[a] = \{b \in A \mid a \sim b\}$ is the equivalence class of a .

Ex 0: $A = \mathbb{Z}$. Let \sim be the "parity" equivalence rel'n:

$a \sim b$ if and only if a and b are both even or both odd (same parity)

Reflexive: a has the same parity as itself

Symmetric: If a and b have the same parity, so do b and a

Transitive: If $a \in b$ have the same parity and so do $b \in c$, both a and c have the same parity as b , and thus as each other

There are two equiv. classes:

$[0] = \{\text{even numbers}\}$ and $[1] = \{\text{odd numbers}\}$

Note that $\dots = [-2] = [0] = [2] = [4] = \dots$
representatives

Ex 1: $A = \mathbb{Z}$

$a \sim b$ means $a = b$ or $a = -b$

Reflexive: $a = a$

Symmetric: If $a = \pm b$, $b = \pm a$

Transitive: If $a = \pm b$, $b = \pm c$, then $a = \pm c$

Equivalence classes:

$$[0] = \{0\}$$

$$[1] = \{-1, 1\}$$

$$[2] = \{-2, -2\}$$

\vdots

Ex 7: $A = \mathbb{Z}$

$a R b$ if $a - b$ is 0, 1, or -1

Reflexive, symmetric, but not transitive

eg. $2 R 3$, $3 R 4$, but $2 \not R 4$

not an equiv. rel'n

Many (but not all) equiv. rel's are of the form:

$a \sim b$ means a and b share the same value of _____

Ex 0: parity

Ex 1: abs. value

Ex 4: $A = \{\text{binary strings}\} = \{\emptyset, 0, 1, 00, 01, \dots\}$

$a \sim b$ if a and b have the same length ✓ equiv. rel'n

$[\emptyset] = \{\emptyset\}$ length 0

$[0] = [1] = \{0, 1\}$ length 1

$[00] = \{00, 01, 10, 11\}$ length 2

$[000] = \{000, 001, \dots, 111\}$ length 3

need to show explicitly

Class activity: Determine whether these are equiv. rel's ($A = \mathbb{Z}$)

a) $a \sim b$ if $a|b$

b) $a \sim b$ if $a \leq b$

c) $a \sim b$ if $a = b$

d) $a \sim b$ if $a - b$ is a mult. of 10

e) $a \sim b$ if $a - b$ is a mult. of 17

Every equivalence rel'n corresponds to a set partition
(and vice-versa)

Def: A set partition of A is a set of subsets
 A_1, A_2, \dots s.t.

$$A_i \cap A_j = \emptyset \quad \text{and} \quad A_1 \cup A_2 \cup \dots = A$$

i.e. every elt. of A is in exactly one A_i

The A_i correspond to the equiv. classes of an equiv. rel'n.

equiv. rel'n \longleftrightarrow equiv. classes $\stackrel{=}{\equiv}$ set partition

Ex 4 (cont.): The set partition corresp. to this
equiv. rel'n is

$$A = A_0 \cup A_1 \cup \dots$$

where

$$A_i = \{\text{strings of length } i\}$$

Ex 15: Let $A = \{\text{binary strings of length 12}\}$.

Set partition

$$\left. \begin{array}{l} A_{000} = \{\text{strings starting w/ } 000\} \\ A_{001} = \{\text{" " w/ } 001\} \\ \vdots \\ A_{111} = \{\text{" " w/ } 111\} \end{array} \right\} \begin{array}{l} \text{set partition into} \\ 8 \text{ sets} \end{array}$$

Corresp. equiv. rel'n:

$a \sim b$ if and only if a and b have the same
first 3 digits

Ex 13: Let $A = A_1 \cup A_2 \cup A_3$ be a set partition with

$$A_1 = \{1, 2, 3\} \quad A_2 = \{4, 5\} \quad A_3 = \{6\}$$

Class activity (if time): Find the corresp. equiv. rel'n.