

Class cancelled next Monday (4/8)

## Galois gps. of polys.

Let  $f(x) \in F[x]$ ,  $K = \text{Sp}_F f$

Def: The Galois gp. of  $f(x)$  is  $\text{Gal}(f) := \text{Gal}(K/F)$

We want to understand  $\text{Gal}(f)$  for different polys.

Thm (Abel, Ruffini): The degree-5 poly. is not solvable by radicals

We know: If  $\deg f = n$ ,  $\text{Gal}(f) \leq S_n$

Generic version:

$K = F(\underbrace{x_1, \dots, x_n}_{\text{think of these as "roots" of a "generic poly" }}) = \text{field of fractions of } F[x_1, \dots, x_n] = \left\{ \frac{3x_1^2 x^3 - 5x_2}{1 + x_4 + x_1^4 x_2}, \dots \right\}$

Have  $S_n \leq \text{Aut}(K/F)$  (permute the  $x_i$ 's)

Set  $L = \text{Fix } S_n$ , and we have  $\text{Gal}(K/L) = S_n$   
↖ field of symmetric functions

Example elts:

•  $f \in F$

•  $e_1 = e_1(x_1, \dots, x_n) = x_1 + x_2 + \dots + x_n$

•  $e_2 = \sum_{i < j} x_i x_j = x_1 x_2 + x_1 x_3 + x_2 x_3 + \dots$

⋮

•  $e_k = \sum_{i_1 < \dots < i_k} x_{i_1} \dots x_{i_k}$

elementary sym.  
funcs.

(D&F call them  $s_k$ )

Fun. Thm. of Sym. Funcs:  $L = F(e_1, \dots, e_n)$

Pf: Let  $L' = F(e_1, \dots, e_n)$ . Then  $L' \subseteq L$  and

$[K:L] = |S_n| = n!$ , so we just need to show

that  $[K:L'] \leq n!$ . This follows since  $K$  is the

splitting field of the following deg.  $n$  poly

in  $L'[x]$ :

$$\begin{aligned}
f_{\text{gen}}^{(n)}(x) &= \prod_i (x - x_i) \\
&= x^n - (x_1 + \dots + x_n)x^{n-1} + \dots + (-1)^n x_1 \dots x_n \\
&= x^n - e_1 x^{n-1} + \dots + (-1)^n e_n
\end{aligned}$$

□

Def: The discriminant of  $f(x) \in F[x]$  is

$$D = \prod_{i < j} (\alpha_i - \alpha_j)^2$$

where  $\alpha_i$  are the roots of  $F$  in  $k := \text{Sp}_F(f)$

Prop:  $D = 0 \iff f$  is inseparable.

Prop:  $D \in F$

Pf:  $D$  is sym. in the  $\alpha_i$ , so

$$D \in F(\underbrace{e_1(\alpha_1, \dots, \alpha_n), \dots, e_n(\alpha_1, \dots, \alpha_n)}_{\text{coeffs. of } f}) = F$$

□

E.g.:

$$a) f = f_{\text{gen}}^{(2)}(x) = (x-x_1)(x-x_2)$$

$$\begin{aligned} D &= (x_1 - x_2)^2 = x_1^2 - 2x_1x_2 + x_2^2 \\ &= (x_1 + x_2)^2 - 4x_1x_2 \\ &= e_1^2 - 4e_2 \end{aligned}$$

So if

$$f(x) = x^2 + \underset{-e_1}{b}x + \underset{e_2}{c}, \text{ then } D = b^2 - 4c \quad (!)$$

b) If  $f(x) = x^3 + ax^2 + bx + c$ ,

$$D = a^2b^2 - 4b^3 - 4a^3c - 27c^2 + 18abc$$

Take a sqrt:

$$K = F(\alpha_1, \dots, \alpha_n)$$

$$\sqrt{D} = \prod_{i < j} (\alpha_i - \alpha_j)$$

$$\begin{array}{c} | \\ F(\sqrt{D}) \end{array}$$

$$\begin{array}{c} | \\ F = F(D) \end{array}$$

Assume  $\text{char } F \neq 2$

If  $G := \text{Gal}(K/F) = S_n$

then  $\exists \sigma \in G$  w/  $\sigma(\sqrt{D}) = -\sqrt{D}$ . Thus,  $\sqrt{D} \notin F$

e.g.  $\sigma = (12)$

Recall:  $A_n = \left\{ \begin{array}{l} \text{even perms.} \\ \text{of } 1, \dots, n \end{array} \right\} \leq S_n$   
index 2

Prop:  $G \leq A_n \Leftrightarrow \sqrt{D} \in F$

PF:  $\sigma(\sqrt{D}) = \sqrt{D} \Leftrightarrow \sigma$  is even, so

$G \leq A_n \Leftrightarrow \sigma(\sqrt{D}) = \sqrt{D} \quad \forall \sigma \in G$

$\Leftrightarrow \sqrt{D} \in \text{Fix } G = F$

□

Next time! find some Galois gps:

$n=2$ :  $f(x) = x^2 + bx + c \in F[x]$   $K := S_{p_F} f$   $G := \text{Gal}(K/F)$

If  $f$  red.,  $K = F$ ,  $G = \text{id}$

If  $f$  irred., then  $[K:F] = 2$ ,  $G = \mathbb{Z}/2\mathbb{Z} \cong S_2$

$K = F(\sqrt{D}) = F(\alpha_1 - \alpha_2) = F(\sqrt{b^2 - 4c})$

(Roots are  $\frac{-b \pm \sqrt{b^2 - 4c}}{2}$ )