

Last class w/ new material (Wed. review)

Recall:

Def: (complex) projective space is the set

$$\mathbb{P}^n(\mathbb{C}) = \{ \text{lines thru. origin in } \mathbb{C}^{n+1} \}$$

$$= \left\{ \alpha = (a_0, \dots, a_{n+1}) \in \mathbb{C}^{n+1} \setminus \{0\} \right\} / (\alpha \sim \lambda \alpha, \lambda \in \mathbb{C})$$

$$= \left\{ [a_0 : \dots : a_n] \right\}$$

$$\text{Cor: } \mathbb{P}^n(\mathbb{C}) = \mathbb{C}^n \cup \mathbb{P}^{n-1}(\mathbb{C})$$

$$\begin{matrix} \nwarrow & \nearrow \\ \text{first} & \text{first coord. 0} \\ \text{coord. 1} & \end{matrix}$$

Want to define projective varieties in $\mathbb{P}^n(\mathbb{C})$

$$\text{Let } f(x, y, z) = xy - z$$

$$\text{Then } f(1, 1, 1) = 0$$

$$f(2, 2, 2) = 2$$

So what does $f([1:1:1])$ mean?

Problem: When we scaled the variables, we doubled z but quadrupled xy

Fix:

Def: $f(x_0, \dots, x_n) \in \mathbb{C}^{n+1}$ is homogeneous of degree d if every term has degree d

If f homog. of degree d

$$f(\lambda a_0, \dots, \lambda a_n) = \lambda^d f(a_0, \dots, a_n)$$

If $\lambda \neq 0$, $f(\lambda a_0, \dots, \lambda a_n) = 0 \iff f(a_0, \dots, a_n) = 0$

Def: If $f \in \mathbb{C}[x_0, \dots, x_n]$ homog.,

$$V(f) := \left\{ [a_0 : \dots : a_n] \in \mathbb{P}^n(\mathbb{C}) \mid f(a_0, \dots, a_n) = 0 \right\}$$

is the projective variety assoc. to f .

Note: no ideal consists of only homog. polys.

Write

$$\mathbb{C}[x_0, \dots, x_n] = \bigoplus_{d=0}^{\infty} A_d$$

$$\text{where } A_d = \left\{ f \in \mathbb{C}[x_0, \dots, x_n] \mid f \text{ is homog of deg. } d \right\}$$

Any $f \in \mathbb{C}[x_0, \dots, x_n]$ can be written uniquely as

$$f = f_0 + f_1 + \dots, \quad f_d \in A_d$$

Def: An ideal $I \subseteq \mathbb{C}[x_0, \dots, x_n]$ is homogeneous if

$$f \in I \Rightarrow f_d \in I \quad \forall d$$

This is equiv. to: I has a generating set consisting only of homog. polys.

Ex: $\mathbb{C}[x, y]$

a) $I = (x+y, x^2+y^2)$ is homogeneous
 $= (x+y, x+y+x^2+y^2)$

b) $J = (y-x^2)$ is not homog.

Def: Let $I \subseteq \mathbb{C}[x_0, \dots, x_n]$ be a homog. ideal. Then

$$\begin{aligned} V(I) &= \{ \alpha = [a_0 : \dots : a_n] \in \mathbb{P}^n(\mathbb{C}) \mid f(\alpha) = 0 \quad \forall f \in I \} \\ &= V(f^{(1)}) \cap \dots \cap V(f^{(k)}) \end{aligned}$$

if $f^{(i)}$ homog. and $I = (f^{(1)}, \dots, f^{(k)})$

These $V(I)$ are called projective varieties

Prop: $I(V) := \{ f \in \mathbb{C}[x_0, \dots, x_n] \mid f(\alpha) = 0 \quad \forall \alpha \in V \}$ is a homog. ideal

Prop: If I homog., \sqrt{I} is homog.

Projective Nullstellensatz: \exists inc. reversing inv. bijections

$$\left\{ \begin{array}{l} \text{nonempty} \\ \text{projective} \\ \text{varieties} \end{array} \right\} \xrightleftharpoons[\vee]{I} \left\{ \begin{array}{l} \text{radical homog. ideals} \\ \text{properly cont. in } (x_0, \dots, x_n) \end{array} \right\}$$

For these varieties/ideals, $V(I(V)) = V$ and $I(V(I)) = \sqrt{I}$.

What about ϕ ? any ideal $\subseteq (x_0, \dots, x_n)$

$$I(\phi) = \mathbb{C}[x_0, \dots, x_n] \quad \text{and} \quad V(\mathbb{C}[x_0, \dots, x_n]) = \phi$$

But also:

$$V((x_0, \dots, x_n)) = \{ \text{pts. in } \mathbb{P}^n(\mathbb{C}) \text{ where } x_0 = \dots = x_n = 0 \} = \emptyset$$

So,

$$\begin{array}{ccc} & I & \rightarrow \mathbb{C}[x_0, \dots, x_n] \\ \phi & \xleftarrow{\vee} & \downarrow \\ & V & \\ & \swarrow & (x_0, \dots, x_n) \end{array}$$

Furthermore

$$\left\{ \begin{array}{l} \text{irred.} \\ \text{nonempty} \\ \text{projective} \\ \text{varieties} \end{array} \right\} \xrightleftharpoons[\vee]{I} \left\{ \begin{array}{l} \text{prime homog. ideals} \\ \text{properly cont. in } (x_0, \dots, x_n) \end{array} \right\}$$

$$\left\{ \begin{array}{l} \text{irred.} \\ \text{nonempty} \\ \text{projective} \\ \text{varieties} \end{array} \right\} \xrightleftharpoons[\vee]{I} \left\{ \begin{array}{l} \text{prime homog. ideals} \\ \text{properly cont. in } (x_0, \dots, x_n) \end{array} \right\}$$

$$\left\{ \begin{array}{l} \text{points} \\ a = [a_0 : \dots : a_n] \end{array} \right\} \xrightleftharpoons[\vee]{I} \left\{ \begin{array}{l} \text{maximal ideals} \\ I(a) = \left(\frac{x_i}{a_i} - \frac{x_j}{a_j} \mid 0 \leq i, j \leq n \right) \end{array} \right\}$$

Pf Sketch of proj. Nullstellensatz: Let

$$V_i = \{[a_0 : \dots : a_n] \in \mathbb{P}^n(\mathbb{C}) \mid a_i \neq 0\} \cong \mathbb{C}^n$$

$$\text{Then } \mathbb{P}^n(\mathbb{C}) = \bigcup_{0 \leq i \leq n} V_i$$

set $a_i = 1$

lots of overlap

Let I be a homog. ideal properly cont. in (x_0, \dots, x_n) ,
and let $V = V(I)$.

Let $V' = \{a \in \mathbb{C}^{n+1} \mid f(a) = 0 \ \forall f \in I\}$

By the affine Nullstellensatz, $I(V') = \sqrt{I}$

We have

$$(a_0, \dots, a_n) \in V' \setminus \{0\} \Leftrightarrow [a_0 : \dots : a_n] \in V,$$

$$\text{so } \sqrt{I} = I(V') \subseteq I(V)$$

Conversely, if f homog., nonconstant, $f(0) = 0$, so

$$f \in I(V) \Rightarrow f(a) = 0 \ \forall a \in V$$

$$\text{homog.} \quad \Rightarrow f(a) = 0 \ \forall a \in V'$$

$$\Rightarrow f \in \sqrt{I}.$$

The rest follows by similar arguments to the affine case. \square