

Last time: Character orthogonality

Today: column orthogonality, examples

Prop 15: The number of irreps. of G equals the number of conj. classes of G .

Pf: By cor. 11, we already know \leq .

For \geq , suppose $\alpha \in \mathbb{C} \text{cl}(G)$ and (α, χ_V) for all irreps V . We will show that $\alpha = 0$, and the result follows.

For any G -repn V , set

$$\varphi_V := \sum_{g \in G} \alpha(g) \rho_V(g) \in \text{End } V$$

One can check that in fact $\varphi_V \in \text{End}_G V$.

[F-H Prop. 2.28]

By Schur's Lemma, $\varphi_V = \lambda \cdot \text{Id}$, so

$$\begin{aligned} \lambda &= \frac{1}{\dim V} \text{Tr}(\varphi_V) = \frac{1}{\dim V} \sum_{g \in G} \alpha(g) \chi_V(g) \\ &= \frac{|G|}{\dim V} (\alpha, \chi_V^*) = 0. \end{aligned}$$

So $0 = \varphi_V = \sum_{g \in G} \alpha(g) P_V(g)$ for any G -repn V .

But if $V = V_{\text{reg}}$, then $\{P_V(g), g \in G\}$ are linearly independent matrices (PF: $P_V(g)v_e = v_g$, and the v_g 's are linearly indep. vectors). Therefore, $\alpha = 0$. \square

Cor 16: The irred. chars. form an orthonormal basis of \mathbb{C} . Furthermore, the columns of the character table are also orthogonal:

$$\sum_{\chi \text{ irred}} \overline{\chi(g)} \chi(h) = \begin{cases} | \text{centralizer of } g |, & \text{if } g=h \\ 0, & \text{otherwise} \end{cases}$$

PF: The first statement follows from Thm. 10 and Prop 15. Column orthogonality is a consequence of squareness and row orthogonality. The details are left as an exercise for the reader (and potentially HW2). \square

Example: S_4

	1	6	8	6	3
	()	(12)	(123)	(1234)	(12)(34)
χ_{triv}	1	1	1	1	1
χ_{sgn}	1	-1	1	-1	1
χ_{ref}	3	1	0	-1	-1
$\chi_{\text{ref}} \otimes \chi_{\text{sgn}}$	3	-1	0	1	-1
χ_W	2	0	-1	0	2

Notes:

$V_{\text{perm}} = \mathbb{C}^4$ $w \cdot v_i := v_{w(i)}$ has character:

$\chi_{\text{perm}}: \quad 4 \quad 2 \quad 1 \quad 0 \quad 0$

decomposes as $\chi_{\text{triv}} + (\text{something})$, and use inner product to show that \wedge is irred.

for W : use column orthogonality

What about A_4 ?

	1 ()	4 (123)	4 (132)	3 (12)(34)
χ_{triv}	1	1	1	1
χ_{ref}	3	0	0	-1
χ	1	ρ	ρ^2	1
χ'	1	ρ^2	ρ	1

Notes:

$$\chi_w \quad 2 \quad -1 \quad -1 \quad 2$$

has $(\chi_w, \chi_w) = 2$

use orthog. to compute the characters.

We obtained characters of A_4 by "restricting" characters of S_4 . Wouldn't it be nice to go the other way?

Def 17: Let $H \leq G$.

a) If (ρ, V) is a G -repn, the restriction of (ρ, V) to H is the H -repn.

$$\text{Res}_H^G(\rho, V) = (\rho|_H, V). \quad \left(\begin{array}{l} \text{or } \text{Res}_H^G V \\ \text{or } \text{Res}_H^G \rho \end{array} \right)$$

b) Let $\sigma_1, \dots, \sigma_k$ be a set of representatives for G/H .

If (π, W) is an H -repn, the induced repn of (π, W) to G is the G -repn

$$\text{Ind}_H^G(\pi, W) = (\rho, V)$$

where

$$V := \bigoplus_{i=1}^k W_i$$

and

$$\rho(g)w_i = (\pi(h_i)w)_j$$

where

$$g\sigma_i = \sigma_j h$$

$\begin{matrix} \in & \in \\ G/H & H \end{matrix}$

Ind_H^G doesn't depend on the choice of coset reps. (up to isom.)

Also note that

$$\dim \text{Ind}_W^G W = |G:H| \dim W$$

and

$$\text{Res}_H^G \text{Ind}_H^G W \cong |G:H| W$$

Ex:

a) $V_{\text{reg}} = \text{Ind}_1^G V_{\text{triv}}$

b) More generally,

$$\text{Ind}_H^G V_{\text{triv}}$$

is the permutation repn. (HW1 #3) of the action of

$$G \text{ on } G/H: \sigma h \cdot v_\tau := v_{\sigma\tau}$$

$\begin{matrix} \in & \in \\ G/H & h \end{matrix}$

Next time: characters of induced reps,
and Frobenius reciprocity